



Review Article

Effects of Ginger (*Zingiber officinale*) as a Dietary Supplement on the Physiology and Health of Poultry

Mohammad Abdalrahem Alhussen* 

College of Agricultural Science and Engineering, Hohai University, Nanjing, China

* **Corresponding author:** Mohammad Abdalrahem Alhussen, College of Agricultural Science and Engineering, Hohai University, Nanjing, China. Email: mohammadalhussen97@gmail.com

ARTICLE INFO

Article History:

Received: 23/04/2026

Revised: 20/05/2026

Accepted: 08/06/2026

Published: 30/06/2026



Keywords:

Antioxidant

Gut integrity

Immunity

Poultry

Zingiber officinale

ABSTRACT

The increasing incorporation of antibiotics into poultry feed for growth development and disease prevention has resulted in serious health implications and the emergence of antimicrobial-resistant bacterial strains. The present study aimed to investigate the effects of ginger (*Zingiber officinale*) powder, as a dietary supplement, on immune response, gut integrity, total antioxidant capacity, and hematological and biochemical parameters in different poultry breeds. The present review examined 76 studies on the effects of dietary ginger on the health and physiology of poultry, including broiler and layer chickens, quail, and turkeys. Only peer-reviewed, indexed, full-text articles published in English were considered and investigated. Natural alternatives, such as ginger powder and its derivatives, in poultry production systems have been shown to mitigate antibiotic-related challenges and have been widely incorporated into poultry feed mixtures. Ginger supplementation in the feed of laying hens, broiler chickens, broiler chicks, and quail has enhanced intestinal integrity and increased the immune response against several viral diseases affecting poultry, such as Newcastle disease, infectious bursal disease, and avian influenza. Ginger has reduced glucose, triglycerides, and total cholesterol levels and increased high-density lipoprotein levels, thereby improving hematological parameters such as red blood cell count, hemoglobin, and packed cell volume, as well as kidney and liver functions. Ginger reduced oxidative stress and improved antioxidant status. Dietary ginger supplementation in poultry diets could serve as a natural feed additive to enhance growth, immunity, and overall health across poultry species, although optimal dosage may differ by species, age, and health status.

1. Introduction

Scientific authorities have recently warned against using antibiotics in poultry feed to accelerate growth and combat pathogens, due to associated health and safety risks. These concerns stem primarily from serious health implications, including the development of antibiotic-resistant strains of pathogens¹. Therefore, many studies have been conducted to explore natural alternatives, notably medicinal plants such as ginger, turmeric, and garlic, which contain therapeutic compounds that do not compromise the health of poultry or humans². *Zingiber officinale*, commonly known as ginger, belongs to the family Zingiberaceae and is a perennial herbaceous plant that typically reaches a height of

approximately one meter. Ginger features green, spear-shaped leaves and produces greenish-yellow flowers³. The part of ginger most commonly used is the thick rhizome, which is used globally as a spice and in traditional Chinese medicine⁴. Ginger contains functional aromatic oils and derivatives, including zingiberene, geraniol, citronellyl acetate, terpineol, linalool, borneol, neral, and flavonoids, as well as phenolic compounds^{5,6}. Ginger is widely recognized for its medicinal, anti-inflammatory, and anti-allergenic properties³⁻⁵. In addition to its ability to lower blood glucose levels in broiler chickens⁶, ginger is also noted for enhancing growth, physiological functions, immune responses, and resilience to

Cite this paper as: Alhussen MA. Effects of Ginger (*Zingiber officinale*) as a Dietary Supplement on the Physiology and Health of Poultry. Journal of World's Poultry Science. 2026; 5(2): 28-35. DOI: 10.58803/jwps.v5i2.129



The Author(s). Published by Rovedar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

stress^{7,8}. Given these promising properties, ginger has become a particularly appealing option for substituting antibiotic growth promoters in poultry farming^{7,8}. However, some individual studies have documented positive effects in different poultry species^{8,9}, a comprehensive synthesis of its impacts, particularly on immune response, gut integrity, antioxidant capacity, and hematological and biochemical parameters, has been poorly investigated. The present study aimed to investigate the effects of ginger powder, as a dietary supplement, on immune response, gut integrity, total antioxidant capacity (TAC), and hematological and biochemical parameters in different poultry species, including broiler chickens, laying hens, quail, and turkeys, across different ages.

2. Search criteria

The present study was exclusively based on studies published in indexed journals. Literature searches were conducted utilizing Scopus, Google Scholar, Web of Science, and PubMed. The search was performed using keywords such as *Zingiber officinale*, ginger, poultry, immunity, gut integrity, antioxidants, and hematology. The current investigations focused on studies examining the effects of ginger supplementation on poultry physiology, immunity, gut health, antioxidant status, and hematological parameters in poultry, including broiler chickens, laying hens, quail, and turkeys, at different ages, irrespective of geographic origin. The selected studies were screened and assessed for scientific quality and relevance prior to inclusion in the present study. Only peer-reviewed, indexed full-text articles were included to ensure data quality. Studies lacking complete sections or published in non-indexed journals, as well as non-English articles and irrelevant studies, were excluded. A total of 169 scientific articles were retrieved from electronic databases; after excluding duplicate studies (47), 122 remained. Another 35 studies were excluded after verifying their titles and summaries to ensure they were not relevant to the topic. The remaining studies (87 studies) were read and examined, and 11 studies were excluded for not meeting the required criteria for the present study. Therefore, 76 studies were included in the present study. A total of 37 original studies were selected for data extraction and summary tables; the remaining 39 articles were used to provide theoretical background and support the discussion and interpretation of the findings.

3. Effects of ginger on the physiology and health of poultry

Ginger can be used as a dietary supplement for poultry in powder form, as an extract, or in combination with

other nutritional supplements, either by incorporating it into the diet or by dissolving it in drinking water^{3,4}. Supplementation with ginger in poultry feed has been reported to improve hematological parameters such as hemoglobin and hematocrit¹⁰, enhance immune response¹¹, reduce cholesterol levels¹², increase TAC¹³, and promote kidney and liver function¹⁴. Furthermore, ginger helps prevent diseases and microbial infections, such as salmonellosis caused by *Salmonella* spp., colibacillosis caused by *Escherichia coli* (*E. coli*), and coccidiosis caused by *Coccidia*¹⁵. Poultry products can be free of harmful substances and safe for human health by incorporating ginger into poultry feed mixtures. [Table 1](#) shows the nutrient content of the ginger plant¹⁶.

Table 1. Nutrient content of the ginger

Elements	Content
Moisture	11 ± 1.41 (%)
Dry matter	90 ± 1.41 (%)
Ash	0.66 ± 0.2 (%)
Fiber	13.5 (%)
Fat	4.2 ± 0.2 (%)
Carbohydrates	92 ± 2.12 (%)
Protein	14.69 (%)
Energy	459.56 (Kcal/100 g)
Lead	0.005 (µg/g)
Cadmium	0.002 (µg/g)
Potassium	10245 (µg/g)
Calcium	351 (µg/g)
Sodium	385 (µg/g)
Iron	122 (µg/g)
Zinc	45.9 (µg/g)
Manganese	6.24 (µg/g)
Selenium	0.189 (µg/g)
Nickel	0.12 (µg/g)
Copper	0.018 (µg/g)
Cobalt	0.015 (µg/g)

Source: Al-Hadad et al.¹⁶

3.1. Immunity

Poultry's immune status can be improved by using ginger in its different forms, such as dried ginger powder, ginger extract, and ginger essential oil, or by adding it to several medicinal plant supplements, which in turn increase antibody titers (immunoglobulin levels) against poultry diseases such as coccidiosis, *Salmonella* infections, and Newcastle disease^{17,18}. [Table 2](#) shows the different effects of ginger on the immune status of poultry.

Table 2. Effects of ginger on the immune status of different poultry breeds

Treatment	Poultry species	Effects/outcomes	Reference
Vitamin E, ginger, and L-carnitine at 250 mg/kg, 2 g/kg, and 500 mg/kg of feed	Broiler chickens	An increase in the average antibody titer against the infectious bursal disease	(19)
Red ginger powder at 2% of the feed	Broiler chickens	A decrease in tissue damage in the cecum and ileum, as well as a decrease in the number of <i>Salmonella</i> spp. in the intestine	(20)
Ginseng and ginger powders at 5 g/kg or as a combination of 2.5 g/kg of feed	Broiler chickens	An increase in the number of WBC, the number of RBC, hemoglobin, and the PCV, as well as an increase in the number of total antibodies (immunoglobulins) against the avian influenza virus	(11)
Ginger powder at 0.25, 0.5, and 0.75%	Quail	Increase in immune globulin, WBC, spleen percentage, and bursa size, as well as the number of Harderian glands	(21)
Ginger powder at 125, 250 mg/kg of feed, and bee Propolis at 500, 1000 mg/kg of feed	Quail	Statistically significant increase in the proportion of NDV antibodies against ND, and a statistically significant increase in the number of WBC (lymphocyte count)	(15)
Ginger powder at a rate of 3, 6, 9 g/kg of feed	Broiler chickens	Increase in the level of NDV antibodies in the blood against ND	(22)
The mixture of anise and ginger extracts at 2 g + 4 g, 2.5 g + 5 g, 3 g + 6 g, was added to the drinking water	Broiler chickens	Increase in total antibody titer against ND, bronchitis, and bursa	(23)
Ginger extract at 1 g/kg of feed	Ostrich chicks	Increase in total antibody titers against ND and improve the effectiveness of the ND vaccine	(17)
Ginger and garlic aromatic oils mixture at 0.5, 1, 1.5, and 2 ml/L	Broiler chickens	Significant increase in IgG and IgM levels, boosting the immune response	(24)
Ginger powder at a rate of 6 g/L of water	Broiler chickens	An increase in the number of T-cells (+CD4) and an increase in gene expression for IFN- γ in the cecal epithelium. Furthermore, ginger helped treat intestinal coccidiosis.	(18)
Ginger extract at 5 g/kg of feed	Laying hens	Decrease in serum cytokines, including TNF- α , IL-6, IL-1 β , and elevated IL-10, IgG, and IgA levels	(25)

IgA: Immunoglobulin A, IgG: Immunoglobulin G, IgM: Immunoglobulin M, IFN- γ : Interferon gamma, IL-6: Interleukin 6, IL-1 β : Interleukin 1 beta, IL-10: Interleukin 10, ND: Newcastle disease, NDV antibodies: Anti-Newcastle disease virus antibodies, PCV: Packed cell volume, RBC: Red blood cells, TNF- α : Tumor necrosis factor alpha, WBC: White blood cells

The improvement in the immune status of different poultry types when using ginger and other plant supplements may be related to the many properties of ginger supplements, such as anti-inflammatory, antioxidant, and antimicrobial effects, as well as their ability to provide essential minerals that support immune cells²³. Ginger has been observed to enhance antibody levels against diverse viral illnesses, including Newcastle disease, bronchitis, and Gumboro disease, likely attributable to its active constituents, shogaol and gingerol, which mitigate oxidative stress and bolster the immune system²⁶⁻²⁸. Moreover, ginger administration has been associated with an increase in the weight of immune organs such as the spleen, thymus gland, and bursa, owing to its anti-inflammatory^{5,29} and antimicrobial properties³⁰. Ginger powder supplementation at a rate of 0.4% in the diet of broiler chickens increased total protein, albumin, and globulin associated with the humoral immune response³¹. Ginger significantly increased the number of T-cells (CD4), expression of interferon gamma (IFN- γ), and immune cells in broiler chickens, compared with untreated infected chickens, due to its ability to treat infections and reduce contributing factors to infection, such as tumor necrosis factor (TNF) and interleukin six (IL-6)¹⁸. The use of red

ginger powder at 2% of the feed as a dietary supplement for broiler chickens reduced *Salmonella* spp. count by stimulating mucus production, to prevent pathogenic bacteria from sticking to the inner surface of the gut villi; consequently, preserving the immune organs such as the cecum and ileum²⁰. The use of ginger powder in the quail diet at a rate of 2 g/kg of feed increased white blood cell (WBC) counts and globulin levels, indicating immunity, antibody production, and immunoglobulin secretion, thereby boosting immunity¹⁹. Ginger powder in the turkey poult's diet at a concentration of 4 g/kg of feed increased the number of WBC, which could be due to the physiological activity of turkeys to raise their immunity after taking a large amount of the active substances contained in ginger³². Additionally, Valdez et al.²⁷ and Hassan et al.³³ found that ginger significantly increased WBC counts. Therefore, ginger can improve the immune system in poultry against many diseases.

3.2. Intestinal health

Table 3 shows the effects of ginger on the health of poultry's intestines.

Table 3. Effects of ginger on the intestinal health status in different poultry breeds

Treatment	Poultry species	Effects/outcomes	Reference
Ginger powder was used as a multi-strain probiotic at 0.15, 0.20, and 0.25% in the feed	Broiler chickens	An increase in <i>Lactobacillus</i> bacterial counts in the ileum, thereby improving intestinal health	(34)
Ginger powder at 3, 6, and 9 g/kg of feed	Broiler chickens	Increase in intestinal villus length and width, deeper crypts, and a decrease in total bacterial count	(21)
Basil leaves, garlic powder, and ginger powder were each added separately at a rate of 50 ml/L in drinking water	Broiler chickens	Ginger alone demonstrated no coccidial egg count, whereas all treatments collectively exhibited a significant reduction in coccidial lesions.	(35)
Protease enzyme derived from ginger at 0.01 and 0.02% of diet	Quail	significant increase in intestinal villus height and width, therefore, enhancing nutrient digestion	(36)
Ginger and garlic aromatic oils mixture at 0.5, 1, 1.5, and 2 ml/L	Broiler chickens	A significant decrease in yeasts, molds, bacteria, <i>Escherichia coli</i> , and <i>Salmonella</i> spp. was observed, while lactic acid bacteria were observed.	(24)
Ethanol extract of ginger at 200, 400, and 600 mg/kg of feed	Dawu Golden Phoenix laying hens	Improved the beneficial microflora in the cecum and increased the beneficial intestinal probiotics	(37)
Ginger extract at 5 g/kg of feed	Laying hens	Enhanced intestinal microbiota function and increased <i>Lactobacillus</i> count, thereby inhibiting some pathogens and reducing inflammation.	(25)

Ginger powder may enhance villus length and width, as well as increase crypt depth in the duodenum, using the enzyme protease extracted from ginger; consequently, enhancing villus length, supporting nutrient absorption, improving intestinal health, and improving digestion^{38,39}. This improvement in intestinal morphology has been reported by Asghar et al.²², who found that ginger increased villus height and crypt depth. Crypt cells play key roles in secreting electrolytes. Accordingly, water content increases in the intestine, thereby improving digestion and supporting overall poultry health⁴⁰. Furthermore, dietary supplementation with ginger and turmeric powder has been shown to reduce colonization by pathogenic bacteria, including *Salmonella* spp. and *E. coli*, as well as by the protozoan parasite responsible for coccidiosis⁴¹. The administration of ginger yeast cultures at 10 g/kg of feed was observed to enhance the digestion of nutrients, including crude protein and crude fat, in laying hens⁴². This effect was attributed to the presence of bioactive compounds in ginger, namely shogaol and gingerol, which stimulate 5-HT₃ receptors in intestinal cells⁴³. Furthermore,

ginger facilitates the activation of lipase, sucrase, and protease enzymes, thereby supporting gastrointestinal digestion^{42,44}. In Addition, Zhang et al.⁴⁵ demonstrated that administering *Saccharomyces cerevisiae* cultures to laying hens increased digestive enzyme activity in chyme and reduced plasma toxin concentrations. The use of ginger and cinnamon oils in Japanese quail has improved gut health⁴⁶ due to their roles in enhancing feed digestion and their rich antioxidant content. Beneficial gut bacteria, such as *Lactobacillus* and *Bifidobacterium*, reduce histological damage and support epithelial cell health⁴⁷. These findings are consistent with a previous report showing that dietary supplementation with ginger and cinnamon can protect the intestines against microbial and parasitic infections⁴⁸.

3.3. Antioxidant status

Antioxidant status is considered a significant indicator of the capacity to mitigate oxidative stress. The associated studies and findings regarding the effects of ginger on antioxidant status are summarized in [Table 4](#).

Table 4. Effects of ginger on antioxidant status in different poultry breeds

Treatment	Poultry species	Effects/outcomes	Reference
Ginger powder at 5, 10, 15 g/kg of feed and thymol at 200, 400 mg/kg of feed	Broiler chickens	A statistically significant decrease in MDA content and GSH levels was observed in the muscles	(49)
Chinese herbal medicine and ginger mixture at 3.32, 10 g/kg of feed	Laying hens and broiler chickens	Significant increase in nitric oxide levels, GSH-Px, and TAC	(13)
Ginger root powder at 1000 mg/kg of ration	Male quail	A reduction in GSH and MDA levels	(50)
Ginger yeast cultures at 5, 10, 20, 40 g/kg of feed	Laying hens	Enhancement in SOD, GSH-Px, POD, and MDA levels	(42)
Ginger powder at 1 g/kg of feed	Aged broiler cocks	An increase in anterior sperm motility and an increase in TAC in semen	(51)
Ginger powder at 5, 10 g/kg of feed	Laying hens	Enhancement in antioxidant status, gizzard and liver health, and a decrease in MDA content	(52)
Ginger and garlic aromatic oils mixture at 0.5, 1, 1.5, and 2 ml/L	Broiler chickens	Enhancement in antioxidant status, an increase in SOD, CAT, TAC, GSH, GSH-Px levels, and a decrease in MDA level	(24)

CAT: Catalase, GSH: Glutathione, GSH-Px: Glutathione peroxidase, MDA: Malondialdehyde, POD: Peroxidase, SOD: Superoxide dismutase, TAC: Total antioxidant capacity

Poultry growth is hindered by oxidative stress caused by free radicals. As a result, chickens were susceptible to various issues such as dropsy and sudden death syndrome⁵³. Medicinal plants containing antioxidants can be used as nutritional supplements in poultry feed mixtures to reduce oxidative stress⁵⁴. Markers such as malondialdehyde (MDA), TAC, superoxide dismutase (SOD), glutathione (GSH), glutathione peroxidase (GPx), and catalase (CAT) indicate antioxidant status and oxidative stress⁵⁵. Ginger supplementation in quail feed has lowered MDA levels⁵¹, likely because ginger contains gingerol and shogaol compounds that protect cells from oxidative damage caused by free radicals⁵⁶. Additionally, the reduction in serum MDA levels may result from ginger's ability to decrease lipid peroxide formation, thereby enhancing antioxidant activity. Ogbuwu et al.⁵⁷ observed that supplementing the diet of broiler breeder hens with ginger at a concentration of 2.5 g/kg of feed resulted in a reduction in MDA levels compared to the control group.

Ginger and cinnamon oils in quail feed increased TAC, SOD, GSH, GPx, and CAT levels and decreased MDA levels due to their capacity to scavenge free radicals and increase antioxidant enzyme activity⁴⁶. Therefore, ginger is considered a powerful antioxidant⁵⁸. The use of ginger and Chinese herbs as dietary supplements in laying hens has led to increased GSH levels, which may be attributed to their ability to promote GSH capture of free radicals¹⁴. Glutathione, known as the main antioxidant in the body, protects DNA from oxidative damage and neutralizes free radicals. Therefore, ginger is considered necessary to reduce reactive oxygen species that impede cellular regulation⁵⁵.

3.4. Hematological and biochemical parameters

Hematological parameters are important indicators of physiological and health status, and the effects of ginger on these parameters are summarized in [Table 5](#).

Table 5. Effects of ginger on the hematological and biochemical parameters in different poultry breeds

Treatment	Poultry species	Effects/outcomes	Reference
Ginger powder at 2, 4 g/kg of feed	Turkey	Significant increase in RBC, serum albumin, and improved liver enzyme activity, reflecting the positive effect of ginger on blood and liver functions	(32)
Ginger powder at 2, 4, 6 g/kg of feed	Broiler chickens	Significant increase in WBC count, the total protein levels, along with reductions in cholesterol, HDL, VLDL, and triglyceride levels in the blood serum	(41)
Ginger extract at 100 g/ton of feed	Laying hens	Decrease in cholesterol, MDA content in yolk, along with reduced ALT and AST enzyme activity in the blood serum	(59)
Ginger at 0.5%, black pepper at 0.5%, 0.25% ginger + 0.25% black pepper	Broiler chickens	Significant differences in increased PCV and RBC, along with a decrease in ALT, AST, total cholesterol, LDL, and VLDL	(60)
Enzyme phytoprotease derived from ginger at 50 and 100 mg/kg of feed	Broiler chickens	Reduction in triglycerides and LDL levels	(39)
Ginger powder at 0.2, 0.4, 0.6% of feed	Broiler chickens	Significant decrease in triglycerides, total cholesterol, and VLDL and LDL, along with an increase in HDL levels	(10)
Ginger powder at 5, 10, 15 g/kg of feed	Broiler chickens	Increased WBC count and heterophils, improved digestion of dry matter, protein, and crude fiber, along with reduced cholesterol, triglycerides, and VLDL in blood	(33)
Ginger powder at 1 g/kg of feed	Aged broiler roosters	Increased anterior sperm motility and sperm production index in testes, along with elevated TAC in semen, and decreased total protein in blood	(51)
Ginger powder at 15 g/kg of feed	Male quail	Increase in ejaculation volume, sperm concentration, motility, vitality, and egg penetration ability, along with a reduction in total cholesterol level	(61)
Ginger powder at 1000 mg/kg of feed	Male quail	Reduction in blood glucose, cholesterol, and triglyceride levels	(62)
Stevia and ginger extracts at 0.5, 1, and 1.5% of feed	Quail	Reduction in AST, glucose, and cholesterol levels in blood serum	(12)
Ginger oil at 0, 0.5, 1 ml/kg of feed	Quail	Significant differences in urea, cholesterol, and AST levels	(46)
Ginger powder at 0.5% to 1% of feed	Broiler chickens	Decrease in triglycerides, cholesterol, and ALT and AST levels	(14)
Ginger and garlic powder at 0.25%, 0.5%, and 1.25% for each, either alone or in combination	Broiler chickens	No significant differences in PCV, hemoglobin, WBC, and their differential components, with a notable increase in RBC count	(63)
Turmeric at 0.75% + ginger 0.75% of feed	Laying hens	Significant differences in RBC, hemoglobin level, and WBC	(64)

ALT: Alanine aminotransferase, AST: Aspartate aminotransferase, HDL: High-density lipoprotein, LDL: Low-density lipoprotein, MDA: Malondialdehyde, PCV: Packed cell volume, RBC: Red blood cell, TAC: Total antioxidant capacity, VLDL: Very low-density lipoprotein, WBC: White blood cell

Ginger powder in Japanese quail feed was associated with a decrease in blood glucose⁶². This decrease in blood glucose levels can be attributed to the phenolic compounds in ginger, which exhibit hypoglycemic properties⁶⁵. Alternatively, a decrease in blood glucose may be due to the ability of ginger's bioactive constituents to increase insulin secretion, thereby promoting glucose uptake by cells and reducing blood glucose levels⁶⁶. Umoru and Agbaye⁶⁷ demonstrated that ginger reduced blood glucose in broiler chickens. Ginger supplementation in the feed of laying hens has reduced blood triglyceride levels, which may be due to ginger's high fiber content, thereby reducing fat absorption in the intestines⁶⁸, or to the presence of HMG-CoA reductase in ginger enzyme⁶⁹. Ginger contains oleoresin, including shogaol and gingerol⁷⁰, which prevents the absorption of bile salts in the intestines and stimulates the liver to produce new bile salts from cholesterol; consequently, cholesterol levels decrease in blood serum³³. Moreover, it was found that ginger contains compounds that reduce cholesterol absorption from the intestine in laying quail⁷¹. Ginger also lowers blood cholesterol levels by stimulating enzymes involved in glycogen synthesis⁷². The enzymes aspartate aminotransferase (AST) and alanine aminotransferase (ALT), which are produced in the liver, are routinely measured to evaluate hepatic health. When a liver injury occurs, ALT and AST levels are elevated in blood serum⁷³⁻⁷⁵. Ginger powder in the feed of laying hens and broiler chickens was associated with decreased AST and ALT levels in the blood⁷¹. This can be attributed to the presence of gingerols (6-gingerols) in ginger, which protect the liver from

damage⁷⁶. There are criteria for kidney health, namely creatinine and uric acid. The use of aromatic ginger and garlic oils in broiler chickens' feed resulted in decreased creatinine and uric acid levels, owing to the antioxidant and anti-inflammatory properties of these oils²⁴. In contrast, Liu et al.⁴² indicated that using ginger as a feed additive in poultry did not significantly affect AST and ALT levels.

4. Conclusion

The present study demonstrated that dietary supplementation with ginger in poultry significantly improved TAC and enhanced gut integrity. It is noted that ginger supported immune function and regulated key hematological and biochemical parameters, including AST, ALT, cholesterol, glucose, and triglycerides, in different poultry breeds. Adding ginger to poultry diets could improve performance, antioxidant capacity, and birds' overall health; however, more investigation is needed to determine optimal inclusion levels, supplementation duration, and species- and production-system-specific responses, especially under diverse environmental and physiological stress conditions.

Declarations

Authors' contributions

Mohammad Abdalrahem Alhussen conceived the review topic, conducted the literature search, selected relevant studies, analyzed and synthesized the findings, and wrote

and revised the manuscript. The author read and approved the final edition of the manuscript before publication in the present journal.

Availability of data and materials

All datasets are available upon request from the corresponding author. No data were generated in the present study, and all information and findings are derived from previously published studies for this review article.

Competing interests

The author declared no competing interests.

Ethical considerations

The author confirmed that ethical concerns, such as plagiarism, permission to publish, research misconduct, data fabrication or falsification, duplicate submissions, and redundant publication, have been thoroughly reviewed. No AI-assisted technologies were used in the generation of this manuscript.

Funding

The present study received no financial support.

References

- Halder N, Sunder J, De AK, Bhattacharya D, and Joardar SN. Probiotics in poultry: A comprehensive review. *J Basic Appl Zool.* 2024; 85(1): 23. DOI: [10.1186/s41936-024-00379-5](https://doi.org/10.1186/s41936-024-00379-5)
- Urban J, Kareem K, Matuszewski A, Bien D, Ciborowska P, Lutostanski K, et al. Enhancing broiler chicken health and performance: The impact of photobiotic on growth, gut microbiota, antioxidants, and immunity. *Phytochem Rev.* 2025; 24: 2131-2145. DOI: [10.1007/s11101-024-09994-0](https://doi.org/10.1007/s11101-024-09994-0)
- Lai W, Yang S, Lin X, Zhang X, Huang Y, Zhou J, et al. *Zingiber officinale*: A systematic review of botany, phytochemistry and pharmacology of gut microbiota-related gastrointestinal benefits. *Am J Chin Med.* 2022; 50(4): 1007-1042. DOI: [10.1142/S0192415X22500410](https://doi.org/10.1142/S0192415X22500410)
- Deng M, Yun X, Ren S, Qing Z, and Luo F. Plants of the genus *Zingiber*: A review of their ethnomedicine, phytochemistry and pharmacology. *Molecules.* 2022; 27(9): 1-25. DOI: [10.3390/molecules27092826](https://doi.org/10.3390/molecules27092826)
- Srinivasan K. Ginger rhizomes (*Zingiber officinale*): A spice with multiple health beneficial potentials. *Pharma Nutrition.* 2017; 5(1): 18-28. DOI: [10.1016/j.phanu.2017.01.001](https://doi.org/10.1016/j.phanu.2017.01.001)
- Shirooye P, Mokaberinejad R, Ara L, and Moghadam MH. Volatile constituents of ginger oil prepared according to Iranian traditional medicine and conventional method: A comparative study. *Afr J Tradit Complement Altern Med.* 2016; 13(6): 68-73. DOI: [10.21010/ajtcam.v13i6.11](https://doi.org/10.21010/ajtcam.v13i6.11)
- Mohamed AB, Al-Rubaei MAM, and Jalil AQ. Effect of ginger (*Zingiber officinale*) on performance and blood serum parameters of broiler. 2012; 11(2): 143-146. DOI: [10.3923/ijps.2012.143.146](https://doi.org/10.3923/ijps.2012.143.146)
- Qorbanpour M, Fahim T, Javandel F, Nosrati M, Paz E, Seidavi A, et al. Effect of dietary (*Zingiber officinale*) and multi-strain probiotic on growth and carcass traits, blood biochemistry, immune responses and intestinal microflora in broiler chickens. *Animals.* 2018; 8(7): 117. DOI: [10.3390/ani8070117](https://doi.org/10.3390/ani8070117)
- Tasya SA, Nurgartiningasih VMA, Permata FS, Djati MS, Nuningtyas YF, Hamiyanti AA, et al. Effects of ginger, turmeric, teak leaf extracts combined with a probiotic on broiler chicken performance. *World Vet J.* 2025; 15(3): 749-755. DOI: [10.54203/scil.2025.wvj76](https://doi.org/10.54203/scil.2025.wvj76)
- Kairalla MA, Aburas AA, and Alshelmani MF. Effect of diet supplemented with graded levels of ginger (*Zingiber officinale*) powder on growth performance, hematological parameters, and serum lipids of broiler chickens. *Arch Razi Inst.* 2022; 77(6): 2089-2095. DOI: [10.22092/ARI.2022.359958.2524](https://doi.org/10.22092/ARI.2022.359958.2524)
- Hussein VA, and Abdulwahid MT. Impact of dietary ginger and ginseng powder in broiler diet on the physiological traits and immune response of avian influenza. *Indian J Forensic Med Toxicol.* 2022; 16(2): 425-432. DOI: [10.37506/ijfmt.v16i2.18014](https://doi.org/10.37506/ijfmt.v16i2.18014)
- Bayril T, Akdemir F, Gürgöze S, and Orhan C. Effect of dietary stevia and ginger extracts on laying performance, fertility, hatchability, and serum biochemical parameters in laying Japanese quails exposed to heat stress. *J Anim Feed Sci.* 2023; 32(3): 289-296. DOI: [10.22358/jafs/160328/2023](https://doi.org/10.22358/jafs/160328/2023)
- Ibtisham F, Nawab A, Niu Y, Wang Z, Wu J, Xiao M, et al. The effect of ginger powder and Chinese herbal medicine on production performance, serum metabolites and antioxidant status of laying hens under heat-stress condition. *J Therm Biol.* 2019; 81: 20-24. DOI: [10.1016/j.jtherbio.2019.02.002](https://doi.org/10.1016/j.jtherbio.2019.02.002)
- Bouchama C, Yousfi L, El Akhdari S, Chadli N, Samouh K, Raoui S, et al. Effect of feed supplementation with *Curcuma longa* and *Zingiber officinale* spices on the zootechnical, biochemical, and microbiological parameters of broiler chickens. *Not Sci Biol.* 2024; 16(3): 11948. DOI: [10.15835/nsb16311948](https://doi.org/10.15835/nsb16311948)
- Abdulhamid ABA, Alraeboub BM, Mohamed RD, and Qareerah AS. Impact of ginger and propolis-enriched diets on the immune system of Japanese quails: An in-depth analysis. *Afro-Asian J Sci Res.* 2023; 1(4): 241-253. Available at: <https://search.emarefa.net/en/detail/BIM-1684389-impact-of-ginger-and-propolis-enriched-diets-on-the-immune-s>
- Al-Hadad MA, Al-Samarrai RR, and Ahumaih MJ. Phytochemical constituents and nutrient evaluation of *Zingiber officinale*. *Tikrit J Pure Sci.* 2023; 22(3): 89-96. DOI: [10.25130/tjps.v22i3.717](https://doi.org/10.25130/tjps.v22i3.717)
- Azadmanesh R, Gholami-Ahangaran M, Karimi-Dehkordi M, and Fathi Hafsanjani E. The immunomodulatory effects of ginger extract on hematological indices and humoral immune response in ostriches vaccinated against Newcastle disease. *Vet Comp Biomed Res.* 2024; 1(2): 72-76. DOI: [10.22103/vcbr.2024.23657.1022](https://doi.org/10.22103/vcbr.2024.23657.1022)
- Saber A, Mawas AS, Khalil AM, Ahmed AI, Shibat Elhamd DM, Ali E, et al. Therapeutic impact of dietary ginger supplementation in chickens experimentally infected with coccidia: Antioxidant, biochemical, and pathological evaluations. *Front Vet Sci.* 2025; 12: 1511759. DOI: [10.3389/fvets.2025.1511759](https://doi.org/10.3389/fvets.2025.1511759)
- Rehman ZU, Chand N, and Khan RU. The effect of vitamin E, L-carnitine, and ginger on production traits, immune response, and antioxidant status in two broiler strains exposed to chronic heat stress. *Environ Sci Pollut Res.* 2017; 24(34): 26851-26858. DOI: [10.1007/s11356-017-0304-8](https://doi.org/10.1007/s11356-017-0304-8)
- Hecrudeati H, Anisa AK, Widiatmoko KD, Alam SSP, Diari IA, Naprila ZH, et al. Effect of red ginger powder (*Zingiber officinale* var. *rubrum*) as a feed additive for starter and finisher broiler chicken to increase immunoglobulin A and immunoglobulin Y expression and to prevent intestinal injury due to *Salmonella enteritidis* infection. *Vet World.* 2022; 15(6): 1506-1514. DOI: [10.14202/vetworld.2022.1506-1514](https://doi.org/10.14202/vetworld.2022.1506-1514)
- El-kashef MMA. Evaluation of using ginger (*Zingiber officinale*) on growth performance, carcass characteristics, blood biochemistry and immune responses of quail birds. *Egypt Poult Sci J.* 2022; 42(2): 199-212. DOI: [10.21608/epsj.2022.249546](https://doi.org/10.21608/epsj.2022.249546)
- Asghar MU, Rahman A, Hayat Z, Rafique MK, Badar IH, Yar MK, et al. Exploration of *Zingiber officinale* effects on growth performance, immunity and gut morphology in broilers. *Braz J Biol.* 2023; 83: e250296. DOI: [10.1590/1519-6984.250296](https://doi.org/10.1590/1519-6984.250296)
- Raziq F, Mushtaq M, Bughio E, Khan MT, Hameed A, Bachaya HA, et al. Dose dependence studies of aniseed (*Pimpinella anisum*) and ginger (*Zingiber officinale*) extract mixture on growth promoting and immunomodulatory effects in broiler chicks. *Pure Appl Biol.* 2023; 12(1): 197-205. DOI: [10.19045/bspab.2023.120021](https://doi.org/10.19045/bspab.2023.120021)
- Alharbi A, Fekry NM, Osailan R, Bahshwan SMA, Al-Ghamdi FA, Al Ghamdi M, et al. Combinatorial effects of ginger and garlic essential oils on broiler chicken performance, digestive functions, immunity, organ function, oxidative stress and defense genes under heat stress. *Pak Vet J.* 2025; 45(4): 1806-1816. DOI: [10.29261/pakvetj/2025.337](https://doi.org/10.29261/pakvetj/2025.337)
- Tong Y, Wang Y, Zhang J, Guo Y, Yuan T, Chen H, et al. Narrative study on the impact of ginger extract on laying performance, egg quality, inflammatory responses, intestinal barrier function, and cecal microbiome and resistance in laying hens. *Poult Sci.* 2025; 104(10): 105448. DOI: [10.1016/j.psj.2025.105448](https://doi.org/10.1016/j.psj.2025.105448)
- Ayustaningwarno F, Anjani G, Ayu AM, and Fogliano V. A critical review of ginger's (*Zingiber officinale*) antioxidant, anti-inflammatory, and immunomodulatory activities. *Front Nutr.* 2024; 11: 1364836. DOI: [10.3389/fnut.2024.1364836](https://doi.org/10.3389/fnut.2024.1364836)

27. Valdez G, Shyur L, Wang SY, and Chen SE. Phytochemicals in ginger, *Origanum vulgare*, and *Syzygium aromaticum* and their potential as a feed additive against *Clostridium perfringens* in broiler production. *Animals*. 2023; 13(23): 3643. DOI: [10.3390/ani13233643](https://doi.org/10.3390/ani13233643)
28. Nawal KS, Mahbuba AGM, and Nidhal AGM. Impact of ginger powder supplementation in broiler diet on the immune status development and small intestine morphology. *Basra J Vet Res*. 2018; 17(2): 99-112. Available at: https://bjvr.uobasrah.edu.iq/article_160400_515eea4d4e45e44b2821d8daa923f54e.pdf
29. Zhang M, Viennois E, Prasad M, Zhang Y, Wang L, Zhang Z, et al. Edible ginger-derived nanoparticles: A novel therapeutic approach for the prevention and treatment of inflammatory bowel disease and colitis-associated cancer. *Biomaterials*. 2016; 101: 321-340. DOI: [10.1016/j.biomaterials.2016.06.018](https://doi.org/10.1016/j.biomaterials.2016.06.018)
30. Hassan A, Youssef I, Abdel-Atty N, and Abdel-Daim A. Effect of thyme, ginger, and their nano-particles on growth performance, carcass characteristics, meat quality and intestinal bacteriology of broiler chickens. *BMC Vet Res*. 2024; 20(1): 269. DOI: [10.1186/s12917-024-04101-z](https://doi.org/10.1186/s12917-024-04101-z)
31. Hassan R, Mosaad GM, and Abd El-wahab HY. Effect of dietary supplemental ginger on broiler performance, carcass characteristics and blood profile. *Int J Vet Sci*. 2019; 2(1): 108-118. DOI: [10.21608/svu.2019.6404.1000](https://doi.org/10.21608/svu.2019.6404.1000)
32. Ajao AM, Oso AO, Lala AO, Olowofeso O, and Adeniran AD. Effect of varying level of ginger powder in the diet of starter turkeys on performance, haematological and serum biochemical indices. *Niger J Anim Prod*. 2018; 45(1): 212-223. DOI: [10.51791/njap.v45i1.330](https://doi.org/10.51791/njap.v45i1.330)
33. Al-Khalafah H, Al-Nasser A, Al-Surrayay T, Sultan H, Al-Attal D, Al-Kandari R, et al. Effect of ginger powder on production performance, antioxidant status, hematological parameters, digestibility, and plasma cholesterol content in broiler chickens. *Animals*. 2022; 12(7): 901. DOI: [10.3390/ani12070901](https://doi.org/10.3390/ani12070901)
34. Qorbani M, Fahim T, Javandel F, Nosrati M, Paz E, Seidavi A, et al. Effect of dietary ginger (*Zingiber officinale* Roscoe) and multi-strain probiotic on growth and carcass traits, blood biochemistry, immune responses and intestinal microflora in broiler chickens. *Animals*. 2018; 8(7): 114. DOI: [10.3390/ani8070114](https://doi.org/10.3390/ani8070114)
35. Okusanya PO, Akinlade OO, Jarikre TA, and Okiya MA. Growth performance, gut histology and anti-coccidial effect of aqueous blends of scent leaf, ginger and garlic in broiler chicken. *Niger Vet J*. 2023; 44(4): 1-10. DOI: [10.4314/nvj.v44i4.2](https://doi.org/10.4314/nvj.v44i4.2)
36. Sultan A, Aziz T, Islam Z, Uzair MS, Alhindary IA, Khan RU, et al. Effect of ginger-based zingibain enzyme on growth and intestinal health in Japanese quails. *Arch Anim Breed*. 2024; 67(4): 571-581. DOI: [10.5194/aab-67-571-2024](https://doi.org/10.5194/aab-67-571-2024)
37. Jin S, Shi J, Zhao M, Liu X, Yang K, Shang E, et al. The influence of dietary supplementation with ginger ethanol extract on laying hens' production performance, antioxidant capacity, and gut microbiota. *Front Vet Sci*. 2025; 12: 1652982. DOI: [10.3389/fvets.2025.1652982](https://doi.org/10.3389/fvets.2025.1652982)
38. Jaric MD, Gottstein Z, Vince S, Zaja IZ, Brus M, Duricic D, et al. Effect of standardized ginger (*Zingiber officinale* Roscoe) extract on gut morphology, microbiota composition, and growth performance in broiler chickens. *Agriculture*. 2025; 15(13): 1448. DOI: [10.3390/agriculture15131448](https://doi.org/10.3390/agriculture15131448)
39. Ahmad U, Sultan A, Khan S, and Tahir M. Effect of ginger derived phyto-protease on production performance, serum biochemistry, nutrient digestibility, gut morphometry and immunity of broilers fed high level of poultry by-product meal-based diet. *Pak J Zool*. 2024; 56(3): 1233-1242. DOI: [10.17582/journal.pjz/20221031071042](https://doi.org/10.17582/journal.pjz/20221031071042)
40. Karangiya VK, Savsani HH, Patil SS, Garg DD, Murthy KS, Ribadiya NK, et al. Effect of dietary supplementation of garlic, ginger, and their combination on feed intake, growth performance, and economics in commercial broilers. *Vet World*. 2016; 9(3): 245-250. DOI: [10.14202/vetworld.2016.245-250](https://doi.org/10.14202/vetworld.2016.245-250)
41. Shewita R, and Taha A. Influence of dietary supplementation of ginger powder at different levels on growth performance, haematological profiles, slaughter traits and gut morphometry of broiler chickens. *S Afr J Anim Sci*. 2018; 48(6): 997-1008. DOI: [10.4314/sajas.v48i6.1](https://doi.org/10.4314/sajas.v48i6.1)
42. Liu J, Jin Y, and Yang J. Influence of spent ginger yeast cultures on the production performance, egg quality, serum composition, and intestinal microbiota of laying hens. *Anim Biosci*. 2022; 35(8): 1205-1214. DOI: [10.5713/ab.21.0514](https://doi.org/10.5713/ab.21.0514)
43. Ali S, Sultan A, Islam Z, Uzair M, Almutairi M, Alrefaei A, et al. Influence of different levels of ginger-derived liquid protease on growth performance, nutrient digestibility, immune organs and gut pH in broiler chickens. *J Anim Physiol Anim Nutr*. 2025; 109(5): 1174-1181. DOI: [10.1111/jpn.14129](https://doi.org/10.1111/jpn.14129)
44. Khan A, Sultan A, Islam Z, Uzair M, Alhindary I, Khan R, et al. Optimising growth performance, nutrients digestibility, immunity and gut health in broilers through ginger-derived phyto-protease enzyme (zingibain) supplementation. *Ital J Anim Sci*. 2024; 23(1): 1695-1703. DOI: [10.1080/1828051X.2024.2416586](https://doi.org/10.1080/1828051X.2024.2416586)
45. Zhang JC, Chen P, Zhang C, Khalil MM, Zhang NY, Qi DS, et al. Yeast culture promotes the production of aged laying hens by improving intestinal digestive enzyme activities and the intestinal health status. *Poult Sci*. 2020; 99(4): 2026-2032. DOI: [10.1016/j.psj.2019.11.017](https://doi.org/10.1016/j.psj.2019.11.017)
46. Abd El-Hack M, Abo ElMaati M, Abusudah W, Awlya O, Almohmadi N, Fouad W, et al. Consequences of dietary cinnamon and ginger oils supplementation on blood biochemical parameters, oxidative status, and tissue histomorphology of growing Japanese quails. *Poult Sci*. 2024; 103(2): 103314. DOI: [10.1016/j.psj.2023.103314](https://doi.org/10.1016/j.psj.2023.103314)
47. Meligy AM, Abd El-Hamid MI, Yonis AE, Elhaddad GY, Abdel-Raheem SM, El-Ghareeb WR, et al. Liposomal encapsulated oregano, cinnamon, and clove oils enhanced the performance, bacterial metabolites antioxidant potential, and intestinal microbiota of broiler chickens. *Poult Sci*. 2023; 102(6): 102683. DOI: [10.1016/j.psj.2023.102683](https://doi.org/10.1016/j.psj.2023.102683)
48. Sahoo N, Mishra S, Swain R, Acharya A, Pattnaik S, Sethy K, et al. Effect of turmeric and ginger supplementation on immunity, antioxidant, liver enzyme activity, gut bacterial load and histopathology of broilers. *Indian J Anim Sci*. 2019; 89: 774-779. DOI: [10.56093/ijans.v89i7.92046](https://doi.org/10.56093/ijans.v89i7.92046)
49. Zidan DE, Kahilo KA, El-Far AH, and Sadek KM. Effects of dietary supplementation of ginger (*Zingiber officinale*) and thymol on the performance, antioxidant status and immune response in broilers. *Glob Vet*. 2016; 16(6): 530-538. DOI: [10.5829/idosi.gv.2016.16.06.10442](https://doi.org/10.5829/idosi.gv.2016.16.06.10442)
50. Abdul-Majeed AF, Abdul-Rahman SY, and Al-krad HA. Effect of ginger on oxidative stress induced by hydrogen peroxide in male quail. *Proceedings of 2nd International Multi-Disciplinary Conference Theme: Integrated Sciences and Technologies, IMDC-IST*. Sakarya, Turkey. 2021. DOI: [10.4108/ea17-9-2021.2315485](https://doi.org/10.4108/ea17-9-2021.2315485)
51. Nemati Z, Dehghani P, Karimi A, Amirdahri S, and Kianifard D. Effects of ginger (*Zingiber officinale*) supplementation on testicular histology, semen characteristic, blood plasma parameters and reproductive performance in aged broiler breeder roosters. *J Anim Physiol Anim Nutr*. 2022; 107(9): 1-13. DOI: [10.1111/jpn.13779](https://doi.org/10.1111/jpn.13779)
52. Derese D, Sun H, Xiong X, Li Z, Malyar R, Lu L, et al. Effects of dietary ginger (*Zingiber officinale*) rhizome powder supplementation on productive performance, egg quality, antioxidant capacity, and hepato-intestinal morphology in pre-peak Xiaoshan laying hens. *Animals*. 2025; 15(15): 2315. DOI: [10.3390/ani15152315](https://doi.org/10.3390/ani15152315)
53. Abubakar JO, Uchechi NC, Abosede OO, and Samuel TO. Role of real phytochemical supplementation to protect cardiac, hepatic, nephrotic, and splenic oxidative stress in broiler chickens. *Transl Anim Sci*. 2023; 7(1): 1-8. DOI: [10.1093/tas/txad106](https://doi.org/10.1093/tas/txad106)
54. Apalowo OO, Minor RC, Adetunji AO, Ekunseitan DA, and Fasina YO. Effect of ginger root extract on intestinal oxidative status and mucosal morphometry in broiler chickens. *Animals*. 2024; 14(7): 1084. DOI: [10.3390/ani14071084](https://doi.org/10.3390/ani14071084)
55. Surai PF, Kochish II, and Fisinin VI. Glutathione peroxidases in poultry biology: Part 1. Classification and mechanisms of action. *World's Poult Sci J*. 2018; 74(2): 185-198. DOI: [10.1017/S0043933918000284](https://doi.org/10.1017/S0043933918000284)
56. Jimoh OA, Ayodele AD, Ojo OA, Okin-Aminu HO, and Olorotimi OJ. Effects of turmeric, ginger, cinnamon, and garlic essential oils on HSP70, NFKB, oxidative DNA damage, inflammatory cytokines, and oxidative markers in broiler chickens. *Transl Anim Sci*. 2024; 8: txae127. DOI: [10.1093/tas/txae127](https://doi.org/10.1093/tas/txae127)
57. Ogbuewu IP, Mbajiorgu CA, and Okoli IC. Antioxidant activity of ginger and its effect on blood chemistry and production physiology of poultry. *Comp Clin Pathol*. 2019; 28(3): 655-660. DOI: [10.1007/s00580-017-2536-x](https://doi.org/10.1007/s00580-017-2536-x)
58. Reda FM, Alagawany M, Mahmoud HK, Mahgoub SA, and Elnesr SS. Use of red pepper oil in quail diets and its effect on performance, carcass measurements, intestinal microbiota, antioxidant indices, immunity and blood constituents. *Animal*. 2020; 14(5): 1025-1033. DOI: [10.1017/S1751731119002891](https://doi.org/10.1017/S1751731119002891)
59. Wen C, Gu Y, Tao Z, Cheng Z, Wang T, and Zhou Y. Effects of ginger extract on laying performance, egg quality, and antioxidant status of laying hens. *Animals*. 2019; 9(11): 857. DOI: [10.3390/ani9110857](https://doi.org/10.3390/ani9110857)

60. Aikpitanyi KU, and Egweh NO. Haematological and biochemical profile of broiler chickens fed diets containing ginger and black pepper additives. Niger J Anim Sci. 2020; 22(2): 114-125. Available at: <https://njas.org.ng/index.php/php/article/view/956>
61. Abdelfattah MG, Hussein MT, Ragab SMM, Abou Khalil NS, and Attaai AH. The effects of ginger (*Zingiber officinale*) roots on the reproductive aspects in male Japanese quails (*Coturnix japonica*). BMC Vet Res. 2023; 19(1): 34. DOI: [10.1186/s12917-023-03576-6](https://doi.org/10.1186/s12917-023-03576-6)
62. Abdul-Majeed A, and Al-Krad H. Influence of ginger as an antioxidant on the physiological performance of male quail stressed by hydrogen peroxide. Mesopotamia J Agric. 2023; 51(1): 141-151. DOI: [10.33899/magri.2023.139269.1224](https://doi.org/10.33899/magri.2023.139269.1224)
63. Iheanacho GC, Ajayi EA, Sholademi EL, Abu OA, Akindoyin MO, Abolude AT, et al. Haematology and serum biochemistry of broiler chickens on dietary supplementation of garlic and ginger powder. Am J Agric Sci Eng Technol. 2024; 8(3): 1-9. DOI: [10.54536/ajaset.v8i3.3335](https://doi.org/10.54536/ajaset.v8i3.3335)
64. Li Z, Kumar M, Irfan M, Khan M, Faran G, Abbas G, et al. Impact of dietary turmeric and ginger powder on performance, serum chemistry and gut microbiota in laying hens. J Appl Poult Res. 2025; 34(4): 100605. DOI: [10.1016/j.japr.2025.100605](https://doi.org/10.1016/j.japr.2025.100605)
65. Shahrajabian MH, Sun W, and Cheng Q. Clinical aspects and health benefits of ginger (*Zingiber officinale*) in both traditional Chinese medicine and modern industry. Acta Agric Scand S B Soil Plant Sci. 2019; 69(6): 546-556. DOI: [10.1080/09064710.2019.1606930](https://doi.org/10.1080/09064710.2019.1606930)
66. Mao QQ, Xu XY, Cao SY, Gan RY, Corke H, Beta T, et al. Bioactive compounds and bioactivities of ginger (*Zingiber officinale* Roscoe). Foods. 2019; 8(6): 185. DOI: [10.3390/foods8060185](https://doi.org/10.3390/foods8060185)
67. Umoru JIA, and Agbaye FP. Performance, serum biochemical and haematological response of broiler chickens fed mixture of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) in diets. Eur J Appl Sci. 2022; 10(2): 596-601. DOI: [10.14738/aivp.102.12216](https://doi.org/10.14738/aivp.102.12216)
68. Gaikwad DS, and Fulpagare YG. Effect of dietary supplementation of ginger and cinnamon on blood profile of broilers. Int J Livest Res. 2020; 10(7): 162-166. DOI: [10.5455/ijlr.20200306072321](https://doi.org/10.5455/ijlr.20200306072321)
69. Malekizadeh M, Moeini MM, and Ghazi S. The effects of different levels of ginger (*Zingiber officinale* Roscoe) and turmeric (*Curcuma longa* Linn) rhizomes powder on some blood metabolites and Production performance characteristics of laying hens. J Agr Sci Tech. 2012; 14: 127-134. DOI:
70. Sharma Y. Ginger (*Zingiber officinale*) an elixir of life: A review. Pharma Innov J. 2017; 6(10): 22-27. Available at: <https://www.thepharmajournal.com/archives/>
71. Herve T, Raphaël KJ, Ferdinand N, Victor Herman N, Willy Marvel NM, Cyril D'Alex T, et al. Effects of ginger (*Zingiber officinale*, Roscoe) essential oil on growth and laying performances, serum metabolites, and egg yolk antioxidant and cholesterol status in laying Japanese quail. J Vet Med. 2019; 2019(1): 7857504. DOI: [10.1155/2019/7857504](https://doi.org/10.1155/2019/7857504)
72. Wen C, Liu Y, Ye Y, Tao Z, Cheng T, and Zhou Y. Effects of gingerols-rich extract of ginger on growth performance, serum metabolites, meat activity of heat-stressed broilers. J Therm Biol. 2020; 89: 102544. DOI: [10.1016/j.jtherbio.2020.102544](https://doi.org/10.1016/j.jtherbio.2020.102544)
73. Yavas I, Adabi SG, and Ceylan N. Liver haemorrhagic syndrome in broilers: its effect on serum parameters, antioxidant capacity, liver enzymes and fatty acid profile of liver. Vet Med Sci. 2025; 11(3): e70372. DOI: [10.1002/vms3.70372](https://doi.org/10.1002/vms3.70372)
74. McGill MR. The past and present of serum aminotransferases and the future of liver injury biomarkers. EXCLI J. 2016; 15: 817-828. DOI: [10.17179/excli2016-800](https://doi.org/10.17179/excli2016-800)
75. Shang HM, Zhao JC, Dong XQ, and Guo Y. Inulin improves the egg production performance and affects the cecum microbiota of laying hens. Int J Biol Macromol. 2020; 155: 1599-1609. DOI: [10.1016/j.ijbiomac.2019.11.137](https://doi.org/10.1016/j.ijbiomac.2019.11.137)
76. Joshi D, Srivastav SK, Belemkar S, and Dixit VA. *Zingiber officinale* and 6-gingerol alleviate liver and kidney dysfunctions and oxidative stress induced by mercuric chloride in male rats: A protective approach. Biomed Pharmacother. 2017; 91: 645-655. DOI: [10.1016/j.biopha.2017.04.108](https://doi.org/10.1016/j.biopha.2017.04.108)